

5

## ACCOMMODATING ZONULAR MINI-BRIDGE IMPLANTS

10

### CROSS-REFERENCE TO RELATED APPLICATIONS

This application is based on U.S. Provisional Application Serial No. 60/223,141 entitled "Surgical Correction of Presbyopia By Synthetic Muscle Bridge Implants (Virtual Reading Glasses)" filed on August 7, 2000, the teachings of which are incorporated herein by reference.

### BACKGROUND OF THE INVENTION

#### Field of the Invention (Technical Field):

The invention relates to an apparatus and method for surgical correction of presbyopia and hyperopia and more particularly to surgical corrections for near vision accommodation of the natural eye lens by the contraction of ciliary muscle/zonule assembly.

#### Background Art:

Normally the human eye loses its ability to focus on nearby objects as individuals reach the age of 40. This condition, known as presbyopia, is due to a progressive loss in the elasticity of the lens of the eye. This is caused by the ciliary muscle which can no longer exert the necessary changes in the lens' shape. Normally the ciliary muscle forces the lens in a rounded shape through the action of zonule fibers on the lens capsule to accommodate for viewing near objects.

The conventional optometric solution to the problem of presbyopia is a prescription of reading glasses or, for individuals who already require glasses to correct other refractive errors such as myopia or astigmatism, a prescription of bifocal or multifocal glasses. This is the reason that Benjamin Franklin invented bifocals in 1760. This century has witnessed a revolution in the surgical treatment of ophthalmic disorders and refractive errors of the human eye. For example, some of these treatments include corneal implantations, cataract extraction, phacoemulsification of the

5 lens, intraocular lens implantation, glaucoma implants to control the intraocular pressure, radial keratotomy, excimer laser ablation of the cornea, trabeculoplasty, iridotomy, virectomy, and the surgical buckle treatment for retinal detachment. The recent surgical solutions for myopia, hyperopia and astigmatism are laser photorefractive keratectomy (PRK), LASIK (laser-assisted in-situ keratomileusis) and  
10 RK or radial keratotomy. Modern techniques for correction of human eye refractive errors are corneal implants (Intacs, Keravision rings), scleral implants (SASI, Schachar Accommodative Scleral Implants) and smart constricting bands equipped with artificial muscle structures to change the eye length.

15       The effective focal length of the human eye must be adjusted to keep the image of the object focused as sharply as possible on the retina. This change in effective focal length is known as accommodation, and is accomplished in the eye by varying the shape of the crystalline lens. This is necessary for the human eye to have clear vision of objects at different distances. Generally speaking, in  
20 unaccommodated normal vision, the curvature of the lens is such that distant objects are sharply imaged on the retina/macula. In the unaccommodated vision, close objects are not sharply focused on the retina/macula and their images lie behind the retinal surface. In order to visualize a near object clearly, the curvature of the crystalline lens is increased, thereby increasing its refractive  
25 power and causing the image of the near object to fall on the retina/macula/fovea region. The change in shape of the crystalline lens is accomplished by the action of ciliary muscle/zonule assembly by which, according to the classical theory of accommodation by German Scientist von Helmholtz (von Helmholtz H., "Über die akkommodation des auges", Albrecht von Graefes Arch Ophthalmol, 1855; 1:1-  
30 89), the radial tension in the lens is reduced allowing the lens to become more convex. In the unaccommodated human eye the lens and its capsule are suspended on the optical axis behind the pupil by a circular assembly of radially directed collagenous fibers called the zonules of Zinn. The inner ends of the zonules are attached to the lens capsule and the outer ends are attached to the  
35 ciliary body. The ciliary body is a muscular constricting ring of tissue located just within the outer supporting structure, or sclera of the eye. The ciliary muscle is relaxed in the unaccommodated eye and therefore assumes its largest inner

09250766-01404  
FOOTNOTES

5 diameter. According to the Helmholtz theory, the relatively large diameter of the  
ciliary body in this unaccommodated condition causes a tension on the zonules  
which in turn pull radially outward on the lens capsule, making it less convex. In  
this state, the refractive power of the lens is relatively low and the eye is focused  
for clear vision of distant objects. When the eye is focused on a near object, the  
10 muscles of the ciliary body contract. This contraction causes the ciliary body to  
move forward and inward, thereby relaxing the outward pull of the zonules on the  
equator of the lens capsule and reducing the zonular tension on the lens. This  
allows the elastic capsule of the lens to contract causing an increase in the  
sphericity or convexity of the lens, resulting in an increase in the optical refraction  
15 power of the lens. Therefore, according to Helmholtz's theory, when an individual  
focused his eye for near vision his ciliary muscle in the eye contracted and  
released tension on the lens zonular ligaments allowing the lens to elastically  
rebound and thicken by itself. In a way the theory asserts that the natural stress-  
free state of the lens is when the ciliary muscle contracts and releases the  
20 tension in the zonules to allow the lens to relax back to its natural state, which is  
more convex and spherical.

Recently, Schachar (Schachar RA, Anderson DA., "The mechanism of  
ciliary muscle function" Ann Ophthalmol. 1995; 27:126-132; and Schachar  
25 RA., "Histology of the ciliary muscle-zonular connections", Ann Ophthalmol. 1996;  
28:70-79.) has proposed a radically different theory of accommodation. The  
fundamental departure, according to Schachar, from the classical Helmholtz  
theory is that the tension in the zonule fibers is constant and that the lens capsule  
itself grows in time, by about 20 microns a year, to finally reduce the tension in  
30 the zonules enough to make the action of the ciliary muscle contraction less  
effective on the lens itself. Schachar further maintains that the hardening of the  
lens with time individually differs for people when they reach the approximate age  
of 40, while getting presbyopic is almost universal for people of 40 years old, and  
thus, is not the only reason behind the phenomenon of presbyopia. By the time  
35 one reaches the age of 40, the lens has increased sufficiently in size so that the  
ciliary muscles can no longer exert tension through the equatorial, anterior, and

09759766 : 01.12.01

5 posterior zonular fibers of the lens when the ciliary muscle contracts for accommodation. For this reason, he maintains, one loses the ability to accommodate at approximately the age of 40 and experience progressive loss of near vision as one gets older. Schachar theorizes that presbyopia could be corrected if the ciliary muscles could be stretched a small amount to allow them  
10 to function on the lens. Relatively recently the Schachar Accommodative Scleral Implant (SASI) has been developed. By making a small incision in the outer coat of the eye (the sclera), one can place a small polymethylmethacrylate (PMMA)) arched stent within this scleral tunnel which will exert a stretching effect on the equatorial lens fibers. Four of these are placed around the eye in the sclera  
15 (approximately 2 mm away from the cornea), and approximately 90 degrees apart. These stents are designed to allow restoration of accommodation. From the Schachar hypothesis of accommodation it could be deduced that if one were able to increase the distance between the lens equator and the ciliary muscle, one could again stretch the zonules and reverse the effect of lens growth and to  
20 narrow this distance.

There are several prior art devices and methods in the form of implants and prostheses for the surgical correction of presbyopia, hyperopia, and myopia.

25 U.S. Patent No. 5,354,331 to Schachar discloses how presbyopia and hyperopia are treated by a method that increases the amplitude of accommodation by increasing the effective working distance of the ciliary muscle in the presbyopic eye. This is accomplished by expanding the sclera in the region of the ciliary body. A relatively rigid band having a diameter slightly  
30 greater than that of the sclera in that region is sutured to the sclera in the region of the ciliary body. The scleral expansion band comprises anterior and posterior rims and a web extending between the rims, the anterior rim having a smaller diameter than the posterior rim.

35 In U.S. Patent No. 5,465,737 to Schachar the teachings are similar to those of the '331 patent, except that the expansion of the sclera is accomplished by

10

15

20

25

30

35

5 and an electromagnetic field receiver adapted to interact with the radiator, both of them exerting an electrostimulation effect on the optic nerve and the retina.

U.S. Patent No. 5,782,894 to Israel discloses a device and method for treating presbyopia by which the ciliary muscles of the eyes are electrically  
10 stimulated when the internal rectus muscles of the eyes are activated in order to focus the eyes on objects within the near field of vision.

U.S. Patent No. 4,961,744 to Kilmer, et al., discloses a surgical apparatus for inserting a plastic, split end, adjusting ring into the stroma of the cornea of the  
15 eye wherein the adjusting ring includes, as a part thereof, a dissecting head to part the stroma and provide a pathway for the adjusting ring as the ring is rotated.

U.S. Patent No. 5,300,118 to Silvestrini, et al., discloses an intrastromal corneal ring (ICR) that is adjustable in thickness and has an elongated, flexible,  
20 preferably transparent or translucent body which forms a circle.

U.S. Patent No. 5,824,086 to Silvestrini discloses a pre-formed intrastromal corneal insert. It is made of a physiologically compatible polymer and may be used to adjust corneal curvature and thereby correct vision abnormalities.  
25

U.S. Patent No. 6,051,023 to Kilmer, et al., discloses a surgical apparatus for inserting a plastic, split end, adjusting ring into the stroma of the cornea of the eye which includes a dissecting head to part the stroma and provide a pathway for the adjusting ring as the ring is rotated.  
30

U.S. Patent No. 5,888,243 to Silvestrini discloses an intrastromal corneal ring housing comprising at least one outer layer of a physiologically compatible polymer having a low modulus of elasticity. The inner portion of the hybrid intrastromal corneal ring may be hollow or may contain one or more  
35 physiologically compatible polymers.

5 U.S. Patent No. 5,766,171 to Silvestrini teaches a device and procedure for the correction of optical abnormalities in a human eye. It involves use of an inventive electrosurgical energy probe with specific physical configurations. The process preferably utilizes a high frequency RF electro-desiccation or ablation device. The procedure involves the initial step of forming at least one access site  
10 allowing access to the corneal volume behind the Bowman's Layer. It preferably is placed in the anterior surface of the cornea through and ending posterior to the Bowman's layer of the eye. The electrosurgical probe is then introduced into the access site and, depending upon the visual abnormality to be corrected, the probe is activated to adjust the volume of the corneal stromal layers through  
15 ablation or desiccation. The shape of the volume desiccated or ablated is dependent upon the aberration to be corrected.

09759766 01201  
20 A number of proposals have been made for changing the focal length of the implanted intraocular lens for cataract patients in response to the natural accommodation mechanism of the eye. One type of adaptive lens comprises an artificial lens whose shape is changed in response to the contraction and expansion of the ciliary muscle. This type of lens is disclosed in U.S. Patent No. 4,254,509 to Tennant, which teaches an eye implant which has an optical lens anteriorly convex and posteriorly planar supported on two diametrically opposed  
25 coplanar feet through two supporting members forming an arch. Each supporting member is unitary with the lens and rooted in one of the feet outside the perimeter of the lens to support the lens with the posterior thereof anterior to the plane of the feet. The lens is formed of a rigid biologically inert material. The supporting members are formed of soft biologically supporting material. The  
30 resulting structure, when fixed into the sclera of the eye, will change as to move the lens anteriorly when forces are applied to the feet upon contraction of the ciliary body.

Similar ideas have been disclosed by U.S. Patent No. 4,842,601 to Smith,  
35 U.S. Patent No. 4,888,012 to Horn, et al., and U.S. Patent No. 4,253,199 to Banko. Two other types of adaptive lenses are described in U.S. Patent No.

5 4,994,082 to Richards, et al., and U.S. Patent No. 5,275,623 to Sarfarazi shows a similar type of compound adaptive lens.

U.S. Patent No. 4,892,543 to Turley describes a compound system comprising a fixed lens having curved posterior and anterior surfaces and a  
10 second component, which is positioned axially posterior of the lens.

U.S. Patent No. 4,790,847 to Woods, U.S. Patent No. 5,152,789 to Willis, and U.S. Patent No. 4,409,691 to Levy describe adaptive lens systems utilizing a simple intraocular lens. These systems have focusing capabilities, which are  
15 achieved by axially shifting the lens in response to normal contraction and expansion of the ciliary muscle resulting from changes in range between the eye and an object under observation. These patents describe similar systems for providing motion of the lens. In each case the ciliary muscle controls zonules, which in turn provide tension to a lens capsule in which the lens system is  
20 mounted. The extremities of the capsule press against a radially compelled, spring-like structure, which also forms a relatively large angle of somewhat, less than 90° with the optical axis of the eye. The lens is positioned on the optical axis. Relaxation of the ciliary muscle releases the radial force and allows the spring to form a more nearly flat shape. When the ciliary muscle contracts, the  
25 pressure on the spring is increased by the action of the lens capsule, the angle between the spring and the optical axis is decreased, and the lens moves axially away from the ciliary muscle.

U.S. Patent No. 4,765,329 to Cumming and Redwitz discusses an  
30 intraocular lens insertion instrument for the implantation of intraocular lenses into the human eyes. The instrument is particularly intended for use in conjunction with the insertion of deformable intraocular lenses through extremely small incisions.

35 U.S. Patent No. 6,051,024 to Cumming teaches intraocular lenses for implanting within natural capsular bags of human eyes that have features on



- 5 distal end portions to prevent movement or sliding thereof relative to fibrosis pockets or tunnels defined about proximally adjacent haptic portions to fixate haptics against dislocation.

U.S. Patent No. 5,578,081 to McDonald presents an artificial lens unit  
10 insertable into a capsular eye lens zone from which a natural lens has been removed, comprising the lens having a light refracting optical portion defining an axis, and consisting of plastic; the unit including haptics for positioning the lens in the capsular zone, the haptics extending at angles relative to a plane normal to the axis and passing through the lens; and the haptics' angles characterized in  
15 that the lens is displaced in the direction of the axis by the haptics in response to eye muscle constriction of the periphery of the capsular zone toward the axis.

U.S. Patent No. 4,786,445 to Portnoy and Ting teaches a method of  
attaching a fixation member to an optic comprising providing the optic with a  
20 cavity opening adjacent the peripheral edge of the optic with a shoulder in the cavity, inserting an inner end portion of the fixation member into the cavity, transmitting laser energy through the optic to the inner end portion of the fixation member to cause the inner end portion to become flowable, and allowing the flowable portion of the fixation member to harden and interlock with the shoulder  
25 to resist withdrawal of the fixation member from the cavity.

U.S. Patent No. 5,843,188 to McDonald teaches a method of providing an artificial lens inserted into the eye between the iris and the natural lens zone. McDonald discloses an artificial lens to be compliant and to have anterior and  
30 posterior surfaces, and haptics extending away from the periphery of the artificial lens. The artificial lens is inserted to extend into position between the iris and the zone, and to cause the haptics to extend into adjacency to the ciliary muscles, and allows the haptics to adhere to the ciliary muscles. Subsequent movement of the ciliary muscles causes movement of the haptics transmitted to effect bodily  
35 movement of the lens in posterior and anterior directions to change the angularity of refraction of light passing through the lens toward the eye retina.

5 U.S. Patent No. 5,476,514 to Cumming, U.S. Patent No. 5,496,366 to Cumming and U.S. Patent No. 5,674,282 to Cumming teach an accommodating intraocular lens to be implanted within the natural capsular bag of a human eye from which the natural lens matrix has been removed through an anterior capsulotomy in the bag circumferentially surrounded by a capsular remnant.

10 During a postoperative healing period following surgery, the anterior capsular remnant fuses to the posterior capsule of the bag by fibrosis about haptics on the implanted lens, and the lens is deflected rearwardly to a distant vision position against the elastic posterior capsule of the bag in which the posterior capsule is stretched rearwardly. After fibrosis is complete, natural brain-induced contraction

15 and relaxation of the ciliary muscle relaxes and stretches the fused remnant and increases and reduces vitreous pressure in the eye to effect vision accommodation by the fused remnant, the posterior capsule, and vitreous pressure. A method of utilizing the intraocular lens in a human eye to provide the eye with accommodation and to enable utilization of a lens with a relatively large

20 optic.

U.S. Patent No. 5,258,025 to Fedorov, Zuev, and Aznabaev presents an optical body, a positioning element, and a supporting element are shaped as an integral unit and have the same radius of curvature that provides for full

25 adherence of the integral unit to an intact natural lens. The distance between the diametrically opposite portions of the supporting element is at least equal to the distance between Zinn's zonules on which the corrective lens rests.

U.S. Patent No. 6,013,101 to Israel teaches an intraocular lens assembly

30 for implantation in a human eye, the eye including a ciliary muscle and zonules controlled by the ciliary muscle, the assembly including an optic, having anterior and posterior surfaces depending from a common edge, at least two, preferably rigid, linkage arms, each being attached to the optic at a first position on the arm thereof, and cooperating with ciliary muscle or the zonules at a second position

35 on the arm, and at least two pivots, one of which is rotatably attached to each respective linkage arm intermediate the first and second positions.

09759766 04201  
F02F099260

5 U.S. Patent No. 5,108,429 to Wiley teaches an adjustable focus lens  
apparatus which includes a transparent lens body having a periphery, an  
attachment device adjacent to the periphery of the lens body for mounting the  
lens apparatus in an eye, and a plurality of micromotor devices spaced equally  
about and connected between the periphery of the lens body and the attachment  
10 means, each of the micromotor devices being responsive to an external control  
signal for selectively changing the position of an associated portion of the lens  
body with respect to the cornea and retina so that the functional power and  
astigmatism of the lens can be appropriately adjusted.

15 It is clear from the above patent literature search that no prior art has  
considered the possibility of accommodating the natural eye lens or intraocular  
lens with mini-bridges placed outside the eye natural capsule and particularly  
within the canal of Hannover, a space between the anterior zonular membrane  
and the posterior zonular membrane, the equatorial zonules and the ciliary  
20 muscle to restore natural eye lens accommodation by the contraction of ciliary  
muscle in near vision situation.

25 **SUMMARY OF THE INVENTION**  
**(DISCLOSURE OF THE INVENTION)**

In accordance with the present invention there is provided a method and  
apparatus for surgical correction of presbyopia and hyperopia by the use of mini-  
bridges for contraction of the eye lens on demand. It also teaches surgical  
30 restoration of near vision accommodation to the natural eye lens by the  
contraction of ciliary muscle/zonule assembly. A circularly distributed assembly  
of directly pressure transmitting, zonule fiber-reinforced active mini-bridges is  
surgically placed between the interior surfaces of the ciliary muscle and the  
exterior equatorial surface of the natural eye lens capsule. Accordingly, the  
35 present invention provides a novel method and apparatus for restoring natural  
accommodation to the natural eye lens utilizing one or more mini-bridges which

5 transmit the circularly distributed contraction force of the ciliary muscle to the  
natural lens capsule. This restores natural near vision accommodation, to correct  
presbyopia and hyperopia on demand. The present invention circularly and  
symmetrically squeezes the eye lens through the lens capsule by means of  
radially distributed pressure transmitting mini-bridges when the ciliary muscle  
10 contracts in close vision situations, in such a way to actively perform near vision  
accommodation by changing the curvature of the eye lens. The surgical  
procedure proposed first uses a ciliary muscle relaxant to create a taut  
configuration for the lens/zonules/ciliary muscle assembly and then employs an  
ultrasonic biomicroscope (UBM) and endoscopic or incisional surgery to place  
15 such mini-bridge implants in and around zonular cavities, such as, preferably the  
canal of Hannover so that the two ends of the mini-bridges span the internal  
surfaces of the ciliary muscle to the exterior surface of the natural eye capsule.  
The surgeon, during this surgical placement of mini-bridges continuously  
observes the microscopic details of the ciliary muscle, zonules, lens capsule and  
20 the Hannover canal.

The composite synthetic muscles to be used for the mini-bridges are either  
passive biocompatible mini-bridges made with polymeric gels such as  
polyacrylamide, polymethylmethacrylate (PMMA), polypropylene or silicone  
25 polymers or a composite of them, electromagnetically or mechanically deployable  
structural mini-bridges, liquid or gas inflatable mini-balloons or biocompatible  
synthetic muscles such as electroactive ionic polymeric artificial muscle  
structures, active materials such as preferably piezocerams, piezopolymers,  
electroactive and eletrostrictive polymers, magnetostrictive materials, liquid  
30 crystal elastomers, and electro or magnetorheological materials suitably  
electroded with biocompatible metals such as gold, platinum and titanium and  
thermally deployable shape memory alloy (SMA) or shape memory polymers  
(SMP) structures. The passive form of polymeric mini-bridges may also be light  
curable to enable the surgeon to implant them in liquid form by incisional tools  
35 and light cure them to a solid form in-situ.

5       A primary object of the present invention is to restore natural near vision accommodation to a presbyopic or hyperopic eye.

Another object of the present invention is to enable presbyopes and hyperopes to become emmetropes (normal vision).

10

Yet another object of the present invention is to enable cataract patients to accommodate their implanted intraocular lenses for near and far vision after surgery by the natural contraction and relaxation of their ciliary muscles.

15

Yet another object of the present invention is to enable presbyopes and hyperopes to see near objects without the use of glasses.

20       A primary advantage of the present invention is that it enables a presbyopic or hyperopic eye to use its own ciliary muscle contraction force to restore natural near vision accommodation.

25       Yet another advantage of the present invention is that it enables cataract patients, after surgical placement of their intraocular lenses, to accommodate or focus on near and far objects by the natural contraction and relaxation of their ciliary muscles.

30

Yet another advantage of the present invention is that it employs a simple surgical procedure for the placement of mini-bridges which involves minimally invasive endoscopic or incisional tools to place the mini-bridge implants.

35

Other objects, advantages and novel features, and further scope of applicability of the present invention will be set forth in part in the detailed description to follow, taken in conjunction with the accompanying drawings, and in part will become apparent to those skilled in the art upon examination of the following, or may be learned by practice of the invention. The objects and advantages of the invention may be realized and attained by means of the

5 instrumentalities and combinations particularly pointed out in the appended claims.

### **BRIEF DESCRIPTION OF THE DRAWINGS**

10

The accompanying drawings, which are incorporated into and form a part of the specification, illustrate several embodiments of the present invention and, together with the description, serve to explain the principles of the invention. The drawings are only for the purpose of illustrating a preferred embodiment of the  
15 invention and are not to be construed as limiting the invention. In the drawings:

Fig. 1(a) shows a side view of the eye lens and its capsule attached circumferentially to the zonular fibers, which are in turn attached to the ciliary muscle in an unaccommodated or relaxed state.

Fig. 1(b) shows the side view of Fig. 1(a) in an accommodated or contracted state.

Fig. 2(a) shows a front view of Fig. 1(a).

Fig. 2(b) shows a front view of Fig. 1(b).

Fig. 3(a) shows a close-up view of the eye lens and its capsule attached circumferentially to the zonular fibers of Fig. 1(a).

Fig. 3(b) shows a close-up side view of the eye lens and its capsule attached circumferentially to the zonular fibers of Fig. 1(b).

Fig. 4(a) shows a close-up front view of the eye lens and its capsule  
35 attached circumferentially to the zonular fibers of Fig. 1(a).

5 Fig. 4(b) shows a close-up front view of the eye lens and its capsule attached circumferentially to the zonular fibers of Fig. 1(b).

Fig. 5(a) shows the pre-surgical configuration of a thermally deployable mini-bridge made up of shape memory alloy (SMA) or shape memory polymer  
10 (SMP).

Fig. 5(b) shows the post surgical configuration of a thermally deployable mini-bridge made up of shape memory alloy (SMA) or shape memory polymer (SMP).

15 Fig. 6(a) shows the pre-surgical configuration of an electrically deployable mini-bridge made up of ionic polymeric artificial muscle materials.

Fig. 6(b) shows the post surgical configuration of an electrically deployable  
20 mini-bridge made up of ionic polymeric artificial muscle materials.

Fig. 7(a) shows the pre-surgical configuration of a mechanically deployable mini-bridge made up of a lazy tong-like (scissor-like) deployable structure that can be endoscopically implanted and deployed within the Hannover canal.

25 Fig. 7(b) shows the post surgical configuration of a mechanically deployable mini-bridge made up of a lazy tong-like (scissor-like) deployable structure that can be endoscopically implanted and deployed within the Hannover canal.

30 Fig. 8(a) shows the pre-surgical configuration of a deployable mini-bridge made up of an electroactively deployable ionic polymeric mini-bridge with bonded electrodes.

5 Fig. 8(b) shows the post surgical configuration of a deployable mini-bridge made up of an electroactively deployable ionic polymeric mini-bridge with bonded electrodes.

Fig. 9(a) shows the pre-surgical configuration of a deployable mini-bridge made up of a polymer hydrogel that can be implanted as a small solid grain in dry  
10 form in the canal of Hannover.

Fig. 9(b) shows the post surgical configuration of a deployable mini-bridge made up of a polymer hydrogel that can be implanted as a small solid grain in dry form in the canal of Hannover.

15 Fig. 10(a) shows the pre-surgical configuration of a deployable mini-bridge made up of a liquid or gas inflatable mini-balloon that can be endoscopically implanted.

20 Fig. 10(b) shows the post surgical configuration of a deployable mini-bridge made up of a liquid or gas inflatable mini-balloon that can be endoscopically implanted.

25 Fig. 11(a) shows the pre-surgical configuration of a softly resilient mini-bridge made up of an elastically deformable and implantable material made with polymethylmethacrylate (PMMA), polypropylene, silicone or other polymeric soft materials.

30 Fig. 11(b) shows the post surgical configuration of a softly resilient mini-bridge made up of an elastically deformable and implantable material made with polymethylmethacrylate (PMMA), polypropylene, silicone or other polymeric soft materials.

35 Fig. 12(a) shows the preferred embodiment for surgical implantation of mini-bridges by means of a syringe or an incisional endoscope.



5 Fig. 12(b) shows the general configuration of the tip of the syringe for  
implantation of the mini-bridges.

Fig. 12(c) shows the preferred incisional endoscope to deploy and implant  
the mini-bridges.

10 Fig. 13 depicts the general configuration for the surgical procedure  
employing an ultrasonic biomicroscope (UBM), a syringe or an incisional  
endoscope and a TV monitor to show the microscopic details of the ciliary/zonule  
assembly.

15 Fig. 14 depicts another configuration for another surgical procedure  
employing a specially designed cannula for the placement of the mini-bridges  
during the normal cataract surgery.

20 **DESCRIPTION OF THE PREFERRED EMBODIMENTS**  
**(BEST MODES FOR CARRYING OUT THE INVENTION)**

25 A novel theory of accommodation and surgical correction of presbyopia  
and hyperopia is first presented, which is a unified theory of accommodation.  
This theory serves as the foundation for the present invention. There is credible  
evidence that Helmholtz was right in claiming that some loss of elasticity and/or  
sclerosis does occur in the lens and the lens capsule with age. In addition, it is  
accepted that the lens itself is ectodermal and grows equatorially at a rate of  
30 approximately 20 microns per year. Consequently, it will be harder for the lens to  
elastically deform back to its more spherical state either naturally or by the  
application of hoop pressure on its capsule. It is also theorized that the zonular  
filaments are never loose and lax because that would render the lens dynamically  
unstable, such as in sports activities requiring rapid postural changes of the head  
35 or the body or in the absence of gravity in space. Therefore, the zonular  
filaments are always under compression like the filaments in a soft hairbrush. If

5 one considers a bundle of soft hairs in a soft hair brush, one can easily observe that although individual hair can not take much compression because they simply buckle or collapse, but collectively as a bundle they can sustain considerable compressive pressures. If a hand is placed on a soft hairbrush one can easily feel the compressive resistance of the hair bundle or hair filaments collectively.

10 This example is similar to the zonular fibers and shows that they are under compression rather than tension when engaged in accommodation. From this theory, an effective surgical procedure to correct presbyopia can be constructed. It is further maintained, that the most relaxed and natural state of the eye is when one is looking at distant objects. In this state the ciliary muscle is relaxed and

15 almost stress-free, the zonules are relaxed and almost stress free, with a small amount of compressive force or hoop pressure applied to them by the ciliary muscle. Most important, is that the lens and the lens capsules are relaxed and almost stress free. Any other state of the eye is an accommodative state in which the ciliary muscle is exerting a hoop stress on the lens capsule through

20 compressive action of the zonular filaments. With progressive hardening and the loss of elasticity of the lens, and its ectodermal growth it will become harder and harder for the ciliary muscle to accommodate by contraction. This is because, on the one hand, the lens and the lens capsule are harder to deform, and on the other hand, the growth of the lens causes the zonular filaments to buckle as they

25 press against the ciliary body. Thus, zonules in this bent and buckled state cannot directly transmit the constricting force of the ciliary muscle to the lens capsule. Referring to the hairbrush example, if the hand on the hairbrush bristles is squeezed down, there is a stronger distributed force or pressure resisting against compression. However, as the downward pressure increases, the

30 hairbrush hair bristles collectively buckle and/or collapse and the compressive force or pressure decreases and it will become easier to further deform the brush bundle. Thus, the best solution is to create direct pressure transmission bridges between the ciliary muscle and the lens capsule so that upon contraction of the ciliary muscle, the hoop pressure is directly transmitted to the anterior region of

35 the lens capsule and thus forces the lens to become more spherical, and accommodate for near vision.

5        The present invention describes elements that operate as force  
transmitting bridges, to transmit the contraction force of ciliary muscles to the lens  
capsule. A circularly distributed assembly of them works together to constrict  
circularly the lens capsule to accomplish the task of accommodation or making  
the lens more spherical. All the user has to do is to look at near objects which  
10 causes the ciliary muscle to contract and thus push the bridges which in turn  
push the capsule. On demand, in this invention, means mental demand because  
as soon as one wants to look at near objects, the ciliary muscle being a smooth  
voluntary muscle, contracts and with the mini-bridges, accentuates natural  
accommodation, making the lens more spherical than without the mini-bridges.

15        Fig. 1(a) depicts a partial side view of the eye lens 1 and its capsule  
(anterior capsule 2' and posterior capsule 2) attached circumferentially to the  
zonular fibers 3, which are in turn attached to the ciliary muscle 4 in an  
unaccommodated or relaxed state in the presence of iris 6, anterior chamber 7,  
20 posterior chamber 8, and cornea 9. The implanted mini-bridges 5, which are  
zonule fiber-reinforced composite materials are bridging and transmitting force  
between the equatorial surface of the lens capsule 2 and the exterior surfaces of  
ciliary muscle 4 and processes 4'. The mini-bridges 5 are preferably composite  
synthetic muscles which are either preferably passive biocompatible mini-bridges  
25 made with water-based polymer hydrogels such as polyacrylamide,  
polymethylmethacrylate (PMMA), polypropylene or silicone polymers or light  
curable polymeric melts or a composite of them, liquid or gas inflatable mini-  
balloons, deployable structural mini-bridges such as deployable lazy tong  
structures, or biocompatible synthetic muscles such as electroactive ionic  
30 polymeric artificial muscle structures, active materials such as preferably  
piezocerams, piezopolymers, electroactive, and eletrostrictive polymers,  
magnetostrictive materials, liquid crystal elastomers, and electro or  
magnetorheological materials suitably electroded with biocompatible metals such  
as gold, platinum, and titanium, and thermally deployable shape memory alloy  
35 (SMA) or shape memory polymers (SMP) structures. The detailed working of  
these elements will be described later in this section. However, it must be

5 mentioned here that all these mini-bridge implants, upon placement in the preferably Hannover canal, work like force transmitting bridges or load bearing walls. Thus, they transmit the contraction force of a ciliary muscle to the equatorial region of the corresponding lens capsule. A circularly distributed assembly of these mini-bridges works together to constrict the lens capsule  
10 circularly to accomplish the task of accommodation or making the lens more convex or spherical. All the user has to do is to want to see near objects, in which case his or her ciliary muscles contract naturally to accommodate for near vision.

15 Fig. 1(b) shows the same partial side view of the eye lens 1 and its capsule (anterior capsule 2' and posterior capsule 2) attached circumferentially to the zonular fibers 3, which are in turn attached to the ciliary muscle 4 in an accommodated or contracted state in the presence of iris 6, anterior chamber 7, posterior chamber 8, and cornea 9. The mini-bridges 5, which are zonule fiber-  
20 reinforced composite materials are bridging and transmitting force between the equatorial surface of the lens capsule 2 and the exterior surfaces of ciliary muscle 4 and processes 4'. Note from Figs. 1(a) and 1(b) that upon contraction of the ciliary muscle 4 in near vision situations, which is caused by the user desire to see near objects, the contraction force is directly transmitted through the mini-  
25 bridges 5 to the equatorial exterior surfaces of the capsule 2, which in turn causes the lens 1 to accommodate in near vision, by becoming more spherical or convex.

Fig. 2(a) is a front view of Fig. 1(a) and shows the eye lens 1 and its  
30 capsule attached circumferentially to the zonular fibers 3, which are in turn attached to the ciliary muscle 4 and in the presence of ciliary processes 4' protruding in to the zonular fibers 3 in an unaccommodated or relaxed state in the presence of ora serrata region 4''. The four mini-bridges 5, which are zonule fiber-reinforced composite materials are bridging and capable of transmitting  
35 force between the equatorial surface of the lens capsule 2 and the exterior surfaces of ciliary muscle 4 and processes 4'. In this relaxed and

5 unaccommodated state, no ciliary muscle contraction force is transmitted to the lens capsule and the eye is in a normal distant vision relaxed state and the ciliary muscle is in a relaxed state.

Fig. 2(b) shows the same front view of Fig. 2(a) in an accommodated or  
10 contracted state in the presence of ora serrata region 4". Note from Figs. 2(a) and 2(b) that upon contraction of the ciliary muscle 4 in near vision situations, the contraction force, which is a circularly distributed hoop force, is directly  
transmitted through the mini-bridges 5 to the equatorial exterior surfaces of the capsule 2, which in turn causes the eye lens 1 to become more convex and to  
15 accommodate in near vision.

Fig. 3(a) is a close-up side view of Fig. 1(a) showing the eye lens 1 and its capsule 2 attached circumferentially to the zonular fibers 3, 3', 3" and 3"', which are in turn attached to the ciliary muscle 4 and in the presence of ciliary  
20 processes 4' protruding in to the zonular fibers 3 in an unaccommodated or relaxed state in the presence of the iris 6, posterior and anterior capsules 2 and 2', respectively and the Hannover canal 9 which is the space between the anterior zonules 3' and posterior zonules 3" and encompassing the equatorial zonules 3'''. In this relaxed and unaccommodated state, no ciliary muscle  
25 contraction force is transmitted to the lens capsule and the eye is in a normal distant vision relaxed state and the ciliary muscle is in a relaxed state.

Fig. 3(b) depicts the same close-up side view as in Fig. 3(a) in an accommodated or contracted state. Note that in this accommodated state, upon  
30 contraction of the ciliary muscle 4 and ciliary processes 4' in near vision situations the contraction force, which is a circularly distributed hoop force, is directly transmitted through the mini-bridges 5 to the equatorial exterior surfaces of the capsule 2, which in turn causes the eye lens 1 to become more convex and to accommodate in near vision.

Fig. 4(a) is a close-up partial front view of Fig. 1(a) showing the eye lens 1 and its capsule 2 attached circumferentially to the zonular fibers 3, which are in turn attached to the ciliary muscle 4 and in the presence of ciliary processes 4' protruding in to the zonular fibers 3 in an unaccommodated or relaxed state. In this relaxed and unaccommodated state, no ciliary muscle contraction force is transmitted to the lens capsule and the eye is in normal distant vision state and the ciliary muscle is in a relaxed state.

Fig. 4(b) depicts the same close-up partial front view as in Fig. 4(a) of the eye lens 1 and its capsule 2 attached circumferentially to the zonular fibers 3, which are in turn attached to the ciliary muscle 4 and in the presence of ciliary processes 4' protruding in to the zonular fibers 3 in an accommodated or contracted state. Note from Figs. 4(a) and 4(b) that upon contraction of the ciliary muscle 4 in near vision situations the contraction force is directly transmitted through the mini-bridges 5 to the equatorial exterior surfaces of the capsule 2, which in turn causes the lens 1 to accommodate in near vision.

Figs. 5, 6, 7, 8, 9, 10, 11, 12 and 13 depict a number of possible materials and configurations for the implantable and deployable mini-bridges 5.

Fig. 5(a) depicts a thermally deployable shape memory alloy (SMA) or shape memory polymer (SMP) mini-bridge implant 5' in a pre-transition state (soft state) which can be cooled by ice water and collapsed in to a thin slender column which can be endoscopically implanted in the Hannover canal. Fig. 5(b) shows the same implanted mini-bridge in activated and deployed configuration 5". This configuration is automatically achieved once the implant thermally equilibrate with body temperature and subsequently thermally deployed 5" with body temperature preferably within the Hannover canal to perform bridging action as well as transmitting force between the exterior surfaces of the ciliary muscle and ciliary processes and the equatorial surface of the eye capsule.

5 Fig. 6(a) depicts a mini-bridge made with electrically deployable ionic polymeric materials **10** in the form of a robotic mini-platform with appropriately attached electrodes **11** that can be transcutaneously deployed via an imbedded induction coil **10'** to perform bridging action in a deployed configuration shown in Fig. 6(b). This embodiment also transmits a force between the exterior surfaces  
10 of the ciliary muscle and ciliary processes and the equatorial surface of the eye lens capsule. The ionic polymeric strips **10** reversibly bend under the action of an electric field.

Fig. 7(a) displays a lazy tong-like (scissor-like) deployable structure in a  
15 collapsed state **10''** that can be endoscopically implanted and subsequently deployed, as shown in Fig. 7(b) within the Hannover canal to perform bridging action as well as transmitting force between the exterior surfaces of the ciliary muscle and ciliary processes and the equatorial surface of the eye lens capsule. This kind of mechanical deployment is performed endoscopically by scissor-like  
20 surgical tools.

Fig. 8(a) depicts an electroactively deployable ionic polymeric mini-bridge in a collapsed state **12** and deployed state **12'**, as shown in Fig. 8(b) with bonded electrodes **14** that can be transcutaneously deployed via an imbedded induction  
25 coil **13** and electroactive polymer strips **10**, to perform bridging action as well as transmitting force between the exterior surfaces of the ciliary muscle and ciliary processes and the equatorial surface of the eye lens capsule. The ionic polymeric strips **10** reversibly bend under the action of an electric field.

30 Fig. 9(a) depicts a polymer hydrogel that can be implanted endoscopically as a small solid grain **15** in dry form in the canal of Hannover. This grain **15** of polymer gel, such as polyacrylamide, will absorb enough aqueous humor and subsequently swell, about 1000 times its initial volume to deploy to a larger structure **15'**, as shown in Fig. 9(b), to perform bridging action as well as  
35 transmitting force between the exterior surfaces of the ciliary muscle and ciliary processes and the equatorial surface of the eye lens capsule. This particular

5 mini-bridge implant is a highly preferred embodiment to perform the force bridging action between the contracted ciliary muscle and the lens capsule. This gel mini bridge upon swelling will be mostly filled with the aqueous humor and thus will be a natural element in the Hannover canal which in fact is a river bed or a canal for the circulation of aqueous humor in the eye. The gel mini-bridge upon swelling  
10 will naturally fill the space between the ciliary muscle and the capsule and will naturally intertwine with the zonular fibers.

Fig. 10(a) depicts a liquid or gas inflatable mini-balloon in a deflated state **16** and inflated state **16'** as shown in Fig. 10(b). This mini balloon can be  
15 endoscopically implanted and subsequently inflated preferably within the Hannover canal to perform bridging action as well as transmitting force between the exterior surfaces of the ciliary muscle and ciliary processes and the equatorial surface of the eye lens capsule. Although the shape of the balloon in Fig. 10 is spherical, different balloon shapes can be utilized, for example cylindrical or  
20 multifingered shapes can be used (not shown).

Figs. 11(a) and 11(b) depict elastically deformable and implantable mini-bridges **5** and **5'** made with polymethylmethacrylate (PMMA), polypropylene, silicone polymer or other polymeric soft materials that can be injected, by means  
25 of a syringe **17** and a needle **19**, in liquid form into the Hannover canal and subsequently enlarged and cured to a soft elastic mini-bridge **5** and **5'**. Light curable polymeric melts are also included.

Fig. 12(a) depicts a partial side view of the configuration of the eye lens **1**,  
30 ciliary muscle **4**, ciliary processes **4'**, zonules **3**, iris **6**, the cornea/limbus region **9'** and generally the posterior chamber **8**. This is a preferred configuration for surgical implantation of mini-bridges by means of a syringe **17**, needle **19** or an incisional endoscope **20**. In this configuration the surgeon will use an ultrasonic biomicroscope to clearly see the ciliary muscle/zonules assembly, the endoscopic  
35 surgical tools and the Hannover canal on a TV monitor, or the like, during surgery, to either place or inject the mini-bridge material in to the Hannover canal



5 by either a syringe **17** filled with the implantable material **18** to inject through the a needle **19** or an incisional endoscope **20**.

Figs. 12(b) and 12(c) display the general configuration of the tip (needle) of the syringe **19'**, **19''** or incisional endoscope **20'** and **20''** to deploy and implant  
10 the mini-bridges **5'** and **5''** of the hydrogel, polymethylmethacrylate (PMMA), polypropylene, silicone polymer, light curable polymeric melts or deployable lazy tong-like mini-structures, respectively.

Fig. 13 depicts the general configuration for the surgical procedure  
15 employing an ultrasonic biomicroscope (UBM) **21** to enable the surgeon to clearly see the ciliary muscle **4**/zonules **3**/Hannover canal **9**, lens **1**, lens capsule **2'**, the iris **6** and endoscopic or incisional tools **19** on a TV monitor **22** while the surgeon places the mini-bridge implants by a syringe or an incisional endoscope **17**.

20 Fig. 14 depicts yet another configuration for an alternative surgical procedure employing a specially designed cannula **23** to enable the surgeon to place the accommodating zonular mini-bridge support system **5** for the eye lens **1** after cataract extraction. In this configuration, the surgeon can clearly see the ciliary muscle **4**/zonules **3**/Hannover canal **9**, eye lens **1**, lens capsule **2'**,  
25 while the surgeon places the mini-bridge implants **5** by a specially designed cannula **23**. The method described is the use of mini-bridges **5** after cataract extraction, with the concurrent use of the foldable intraocular lens **1** for accommodative purposes. The technique involves performing cataract extraction using either clear cornea or scleral tunnel approach. Once the normal  
30 capsulorhexis and phacoemulsification of the nuclear and cortical material is performed, the anterior capsular bag **2'** remains with its normal zonular attachments. The goal is to place the mini-bridges **5** at the position of the anterior capsule **2'**, in the usual case at 3 and 9 o'clock positions on the lens. The placement of the mini-bridges **5** can be performed either before the foldable  
35 intraocular lens **1** is placed in the capsule, or after the lens **1** is placed in the capsule and its haptics rotated to the 3 and 9 o'clock positions. The technique

5 involves retracting the already dilated iris 6 (3 and 9 o'clock) positions, and  
placement of the mini-bridges 5, either in the form of composite liquid form or gel  
form, into the respected positions in the Hannover canal 9, thus bridging the  
ciliary muscle 4 to the anterior capsule 2'. This will involve a specific designed  
cannula 23. The shape of the preferred specific designed cannula is shown in  
10 Fig. 14. As shown the curved end 25 of the cannula 23 lifts the iris 6, while the  
liquid or gel is injected into the respective positions. The final placement of these  
mini-bridges can be determined either by an ultrasonic biomicroscope (UBM) or  
by direct endoscopic means. Although only four mini-bridges are shown, different  
arrangements of bridges can be used, such as three or more symmetrically  
15 spaced mini- bridges around the lens capsule. A single ring shaped mini-bridge  
can also be used (not shown).

Thus, the present invention, creates active pressure-transmitting zonule  
fiber-reinforced mini-bridges or force carrying webs between the interior surfaces  
20 of the ciliary body/muscle/processes and the exterior zonular surfaces of the eye  
lens capsule, circularly squeezing the eye lens capsule when the ciliary muscle  
contracts in close vision situations, in such a way to actively change the curvature  
of the eye lens, bringing its focal point to finally coincide with the retina/macula  
region and thus correct presbyopia and hyperopia on demand.

The surgical procedure proposed first uses a ciliary muscle relaxant to  
create a taut configuration for the lens/zonules/ciliary muscle assembly and then  
employs an ultrasonic biomicroscope (UBM) to enable the surgeon to clearly see,  
on a TV monitor, the ciliary muscle/zonules/Hannover canal, lens capsule, iris,  
30 mini-bridges, surgical tools, and thus be able to perform endoscopic or incisional  
surgery to place such mini-bridge implants in and around zonular cavities, such  
as, preferably the canal of Hannover so that the two ends of the mini-bridges  
span the internal surfaces of the ciliary muscle to the exterior surface of the  
natural eye capsule. In general other materials such as Nylon, Proline, light  
35 curable polymers or Heparin or other biologically active compounds to reduce  
body rejection of the mini-bridges can also be used.

5        Although the invention has been described in detail with particular  
reference to these preferred embodiments, other embodiments can achieve the  
same results. Variations and modifications of the present invention will be  
obvious to those skilled in the art and it is intended to cover in the appended  
claims all such modifications and equivalents. The entire disclosures of all  
10 references, applications, patents, and publications cited above, are hereby  
incorporated by reference.

09359768.011201  
T02T0:99265260